



Original Research Article

Amla possesses antifungal potential against *Alternaria alternata*, *Bipolaris specifera* and *Curvuleria lunata*

Arti Heer¹, Vikas Sharma^{1,*}, Navneet Kour¹, Arvind Badyal¹, Shivangi Sharma²

¹Division of Biochemistry, Faculty of Basic Sciences, Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu, Jammu and Kashmir, India

²Dept. of Chemistry, University of Jammu, Jammu and Kashmir, India



ARTICLE INFO

Article history:

Received 18-05-2022

Accepted 28-06-2022

Available online 20-08-2022

Keywords:

Emblica officinalis
Methanolic extract
Alternaria alternata
Biopolaris specifera
Curvuleria lunata

ABSTRACT

Introduction: The present study was carried out to evaluate antifungal activity of methanolic extract from *Emblica officinalis* leaves against three phytopathogens viz., *Alternaria alternata*, *Bipolaris specifera* and *Curvuleria lunata*.

Materials and Methods: Inhibitory potential of methanolic extract was analyzed by poisoned food technique in which different concentrations of test material were prepared in sterilized potato dextrose agar.

Results: Results of the present study revealed that *Emblica officinalis* has potential activity against the three test pathogens with IC₅₀ value of 0.91±0.01 mg/mL, 1±0.015 mg/mL and 1.1±0.0152 mg/mL against *Bipolaris specifera*, *Alternaria alternata* and *Curvuleria lunata* respectively.

Conclusion: Amla has some potential antifungal activity for the prevention and treatment of fungal infections.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Emblica officinalis known as Indian gooseberry or *amla*, belongs to family Euphorbiaceae, is a medium-sized deciduous tree with gray bark and reddish wood. It is native to tropical southern Asia and possesses very highly characteristic medicinal value. *E. officinalis* plant extracts revealed antibacterial / antifungal¹, antioxidant² and cardio-protective³ properties. *E. officinalis* is highly nutritious and is one of the richest sources of vitamin-C, amino acids and minerals.⁴ It contains several chemical constituents like tannins, alkaloids and phenols.⁵ Pharmacological research reports on amla reveals its analgesic⁶, cardio⁷, gastro⁸, nephron⁹ and anticancer¹⁰ properties. In view of the above, the present research was carried out to evaluate the

antifungal activity of *E. officinalis*.

2. Materials and Methods

Preparation of extracts: Fresh leaves of amla were collected from Herbal Garden, Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu (SKUAST-J). The freshly collected leaves were chopped, shade dried and ground into powdered form. The powdered dried material was then extracted with methanol at room temperature to obtain extracts for bioevaluation.

2.1. Determination of antifungal activity of *E. officinalis* extracts by poisoned food technique

Different concentrations of test component (extract) were prepared in sterilized potato dextrose agar and poured in 9 cm petri plates. After this, 5 mm bit of test fungus was



* Corresponding author.

E-mail address: vikas.skuast@gmail.com (V. Sharma).

Table 1: Growth inhibitory effect of *Emblica officinalis* leaves on fungal strains.

Extract	Conc. (mg/mL)	Phytopathogenic fungi		
		<i>Alternaria alternata</i>	<i>Curvuleria lunata</i>	<i>Bipolaris specifera</i>
Methanolic			Growth Inhibition (%)	
	0.5	37.5	37.5	42.5
	1	51.2	47.5	51.0
	2	68.5	66.25	70
	IC ₅₀	1±0.015	1.1 ±0.0152	0.91±0.01
Amphotericin (positive control)			Growth Inhibition (%)	
	10	48.5	46.20	50.75
	20	65.00	61.00	71.50
	40	83.60	81.60	85.69
	IC ₅₀	9.5±0.1	12.1±0.4	5.7±0.2

Maximum growth inhibition by test material as indicated by IC₅₀ value.

**Fig. 1:** Antifungal IC₅₀(mg/mL).**Fig. 2:** Antifungal activity against *Bipolaris specifera*.**Fig. 3:** Antifungal activity against *Alternaria alternata*.

inoculated in the center of the agar plate (mycelia surface of the bit was placed upside down) followed by incubation of petri plates at 26 °C. The extension diameter (mm) of hyphae from the center to the dish was measured at 24 h interval, till the growth of fungus in the plate without test component (control) reached the edge of the plates. The experiment was repeated thrice and results were expressed as average of three replicates.¹¹

Fungal growth diameter in each plate containing concentrations of test component was determined to calculate per cent growth inhibition.

The antifungal indices was calculated as:

$$2.2. \text{ Antifungal index (\%)} = (1 - D_a/D_b) \times 100$$

1. D_a = Diameter of growth zone in the experiment dish (mm).
2. D_b = Diameter of growth zone in the control (mm).

3. Results and Discussion

Exploitation of antifungal activity of amla leaves revealed that methanolic extract of amla inhibited the growth of the colonies of large number of fungal species. The antimicrobial efficacy of extract of amla was qualitatively assessed on the basis of inhibition zone. To evaluate the antifungal activity of methanolic extract from *amla* leaves, three important phytopathogenic fungi, *Bipolaris specifera*, *Alternaria alternata* and *Curvuleria lunata* were selected. The results of the present study showed that the antifungal activity of amla against three test pathogens with IC₅₀ values were 0.91±0.01 mg/mL, 1±0.015 and 1.1±0.0152 mg/ml respectively (Table 1). The antimicrobial components produced by plants are active against plant pathogens. However, their use is increasingly restricted due to the harmful effects of pesticides on human health and the environment. The use of biological compounds extracted from plants may be an alternative to conventionally used fungicides to control phytopathogenic fungi. The search for antimicrobials from natural sources has received much

attention and efforts have been put in to identify compounds that can act as suitable antimicrobials agent to replace synthetic ones. Phytochemicals derived from plant products serve as a prototype to develop less toxic and more effective medicines in controlling the growth of micro-organism.¹² These compounds have significant therapeutic application against human pathogens including bacteria, fungi or virus. Numerous studies have been conducted with the extracts of various plants, screening antimicrobial activity as well as for the discovery of new antimicrobial compounds.¹³ Therefore, medicinal plants are finding their way into pharmaceuticals, nutraceuticals and food supplements. Further, research has to be conducted to find out the possibility of this medicinally important plant as a potent antimicrobial drug and for other pharmacological properties to develop as cost effective formulation.

4. Source of Funding

None.

5. Conflict of Interest

None.

References

- Hossain MM, Mazumder MM, Hossen K, Tanmy SM, Rashid MJ. In vitro studies on antibacterial and antifungal activities of *Emblica officinalis*. *Int J Pharm Sci Res.* 2012;3(4):1124–7.
- Golechha M, Bhatia J, Arya D. Studies on effects of *Emblica officinalis* (amla) on oxidative stress and cholinergic function in scopolamine induced amnesia in mice. *J Environ Biol.* 2012;33(1):95–100.
- Bhattacharya SK, Bhattacharya A, Sairam K, Ghosal S. Effect of bioactive tannoid principles of *Emblica officinalis* on ischemia-reperfusion-induced oxidative stress in rat heart. *Phytomedicine.* 2002;9(2):171–4.
- Srivastuki KP. Nutritional and health care benefits of Amla. *J Pharmacognosy.* 2012;3(2):141–51.
- Zhang LZ, Zhao WH, Guo YJ, Tu GZ, Lin S, Xin LG. Studies on chemical constituents in fruits of Tibetan medicine *Phyllanthus Emblica*. *Zhongguo Zhong Yao Za Zhi.* 2003;28(10):940–3.
- Sharma SK, James B, Christina AJM. Evaluation of anti-pyretic and analgesic activity of *Emblica officinalis* Gaertn. *J Ethnopharmacol.* 2004;95(2):83–5.
- Baliga MS, Prabhu AN, Prabhu DA, Shivashankara AR, Abraham A, Palatty PL. Antidiabetic and cardioprotective effects of Amla (*Emblica officinalis* Gaertn) and its phytochemicals: Preclinical Observations, Bioactive Food as Dietary Interventions for Diabetes. *Biology.* 2013;1(4):583–600.
- Chatterjee A, Chattopadhyay S, Sandip K. Biphasic Effect of *Phyllanthus emblica* L. Extract on NSAID-Induced Ulcer: An Antioxidative Trail Weaved with Immunomodulatory Effect. *Evid Based Complement Altern Med.* 2011;1(13):146808.
- Yokozawa T, Kim HY, Kim HJ, Sugino H. Amla (*Emblica officinalis* Gaertn.) Attenuates age- related renal dysfunction by oxidative Stress. *J Agricultural Food Chem.* 2007;55(19):7744–52.
- Madhuri S. Effect of ProImmu on estrogen induced uterine cytotoxicity in albino rat; 2008. Available from: https://www.researchgate.net/publication/343414122_Effect_of_ProImmu_on_estrogen_induced_uterine_cytotoxicity_in_albino_rat.
- Guleria S, Tiku AK, Koul A, Gupta S, Singh GA, Razdan VK. Antioxidant and antimicrobial properties of the essential oil and extracts grown in North-Western Himalaya. *Sci World J.* 2013;28:790580.
- Ahmad I, Beg AZ. Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multiple drug resistant human pathogens. *J Ethnopharmacol.* 2000;74(2):113–23.
- Guleria S, Kumar A. Antifungal activity of some Himalayan medicinal plants using direct bioautography. *J Cell Mole Biol.* 2006;5:95–8.

Author biography

Arti Heer, Ph.D Scholar

Vikas Sharma, Associate Professor

Navneet Kour, Ph.D Scholar

Arvind Badyal, Student

Shivangi Sharma, Ph.D Scholar

Cite this article: Heer A, Sharma V, Kour N, Badyal A, Sharma S. Amla possesses antifungal potential against *Alternaria alternata*, *Bipolaris specifera* and *Curvuleria lunata*. *Curr Trends Pharm Pharm Chem* 2022;4(3):123-125.