



## Review Article

## Ethnopharmacological review of boswellia serrata for anticancer activity

Amit Gajanan Nerkar<sup>1,2,3,\*</sup>, Pallavi Gade<sup>1</sup><sup>1</sup>Dept. of Pharmacy, CAYMET's Siddhant College of Pharmacy, Sudumbare, Pune, Maharashtra, India<sup>2</sup>Founder and Director, Ateos Foundation of Science Education and Research, Pune, Maharashtra, India<sup>3</sup>Founder and Director, Carolene Therapeutics, Pvt. Ltd, Aurangabad, Maharashtra, India

## ARTICLE INFO

## Article history:

Received 20-10-2022

Accepted 11-11-2022

Available online 21-11-2022

## Keywords:

Boswellia

Anticancer

Ethnopharmacology

Boswellic acids

Phytochemistry

## ABSTRACT

Boswellic acid is an Ursane group compound belonging to triterpenoids. It is the major constituents of the gum derived from plant *Boswellia serrata* Roxb. It belongs to Family Burseraceae. It is also known as *B. glabra* and commonly known as the Salai guggal, white guggal, Indian olibanum. *Boswellia serrata* is medium-sized tree widely distributed in the India and Africa. *B. Serrata* is highly medicinal as well as economically potential. Currently, it is has the potential use in various herbal and ayurvedic formulations for the treatment of various disorders which include inflammation and other types of disorders. Since ancient and centuries anti-cancer drugs discovered from herbal medicines have been used in clinical practice. The active constituent of such formulations being vincristine, vinblastine and the camptothecene derivatives, topotecan, irinotecan and etoposide. There is current need for herbal and ethnopharmacological therapeutics to develop the novel anti-cancer drugs which is safe with effective mechanism. The new discoveries of developing combined ingredients from effective traditional formula or single ingredient as per the traditional medicine theory should be focused. This new approach will promote the academic research and the industry development of traditional medicine. This ethnopharmacological review is mainly focused on anticancer activity of *Boswellia serrata*.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

## 1. Introduction

Herbal medicinal plants are important part of pharmacognosy, ayurveda, and ethnopharmacology. They have been used since the ancient era for the therapeutics related to mankind and for veterinary purposes. Also accordingly as per the 20<sup>th</sup> century the Herbal medicinal plants should be safe for therapeutics and if single ingredient is used should be as per the standards of FDA.<sup>1</sup> Products obtained from the natural origin have been used for the past centuries and today have great significance in pharmaceutical industry. Most of the products derived from the natural sources are now successfully employed

in pharmaceutical formulations as drugs.<sup>2</sup> Boswellic acid consists of a series of pentacyclic triterpene molecules that are produced by the plant *Boswellia serrata*.<sup>3</sup> The potential applications of Boswellic acid for treatment of cancer have been focused hereIt may help the researchers for further investigation.<sup>4</sup> *Boswellia serrata roxb* is also known as Indian olibanum and is found in the central and northern part of east India especially dry hill forest of Rajasthan, Madhya Pradesh, Gujarat and Bihar etc.<sup>5</sup> The species produced olibanum resin of various qualities, which are commonly known as Salahi white guggul or Indian olibanum or kundru.<sup>6</sup>


\* Corresponding author.

E-mail address: [dragnerkar@gmail.com](mailto:dragnerkar@gmail.com) (A. G. Nerkar).

### 1.1. Taxonomical Classification of *B. serrate*<sup>7</sup>

1. Kingdom: Plantae
2. Clade: Tracheophytes
3. Clade: Angiosperms
4. Clade: Eudicots
5. Clade: Rosids
6. Order: Sapindales
7. Family: Burseraceae
8. Genus: *Boswellia*
9. Species; *B. serrata*
10. Common Name: *Boswellia serrata* Roxb.

### 1.2. Phytochemical profile of *Boswellia Serrata*



**Fig. 1:** Active constituents of *Boswellia Serrata*

The *Boswellia serrata* dried gum or the resinous part contains the active constituents such as monoterpenes, diterpenes, triterpenes, tetracyclic triterpene acids and four major pentacyclic triterpene acids i.e. β-boswellic acid, acetyl-β-boswellic acid, 11-keto-β-boswellic acid and acetyl-11-keto-β-boswellic acid, etc.<sup>8</sup>

## 2. Review on Anticancer Activity of *Boswellia Serrata*

Cancer is a crucial disease and co-associated with genetic make up, and mostly risky in geriatric patients and later stages of the life. With advancement of modern research and chemotherapeutics there has been a revolution of in the development of anti-cancer drugs.<sup>9</sup> BSE (*Boswellia Serrata* Extract) is reported to affect the propagation of growing Breast cancer and brain tumor cells. It is a known inducer of apoptosis further in a study it was observed that the ethanolic extract tested showed cytostatic, cytotoxic and apoptotic abilities against leukaemia and brain tumour cells. It was found to induce apoptosis and act as a potent antiproliferative agent.<sup>10,11</sup> In an investigation with BSE which contained 60% BAs have showed that it apparently inhibited tumors and inflammation in mice. BSE has also reported anti-cancer activity in Ehrlic ascites carcinoma and S-180 tumors, in mice. This might occur by possible mechanism of action of inhibiting the cell proliferation and

proliferation by interfering with biosynthesis of DNA, RNA and proteins.<sup>12</sup> It reduces the tumour cell proliferation and induce apoptosis in several in vitro animal experiments.<sup>13,14</sup> The effectiveness of BSE against peri-tumoral oedema could be increased by improving the bioavailability of AKBA.<sup>15,16</sup>

The *Boswellia Serrata* nanoparticle formulation has shown with significant activity to produce anticancer effect for the treatment of prostate cancer.<sup>17,18</sup> Boswellic acid nanoparticles induced apoptosis DNA fragmentation, outcome of which leads to DNA fragmentation which causes high potential of apoptosis.<sup>19</sup> Suhail et al. demonstrated that *Boswellia sacra* essential oil prepared by hydro distillation has been tumor cell-specific cytotoxicity in several type of cancer cells.<sup>20</sup> Consistent with anti-proliferative and proapoptotic properties in cultured breast cancer cells, *Boswellia sacra* essential oil has been observed to produce cytotoxicity in metastatic breast cancer cell lines resistant to a cancer medicine<sup>21,22</sup> Serum concentration of MMP-7, MMP-9 and EGF, and plasma TGF-β and TNF-α levels were assayed by using ELISA procedure.<sup>23</sup> Immunohistochemistry technique was used to estimate the expression of COX-2 and cyclin D1 in the colon. Colonic β-catenin, K-ras and c- myc gene expression was detected by RT-PCR. In addition, histological study of colon tissue was performed. the Colon cancer group showed significant increase in the studied biochemical markers. In the contrast the treated groups all showed significant reduction in these markers. The Colon cancer group had significantly elevated COX-2 and cyclin D1 expression in colon tissue. In contrast, all treated groups showed a marked decrease in COX-2 and cyclin D1 expression<sup>24</sup> The Cancer group showed significant upregulation of the expression level of β-catenin K-ras and c-myc genes in the colon tissue. While all treated groups exhibited significant downregulation of the expression levels of these genes. Histopathological examination of tissue sections from the colon in the cancer group showed dysplasia and anaplasia in epithelial cell lining of the glandular structure. While treatment with 5-fluorouraci extract or *Boswellia serrata* extract showed a marked improvement in the histological characteristics of colon tissue.<sup>25</sup> It was also observed that acetyl-keto-boswellic acid [AKBA] inhibited cell growth in a several colon cancer cell lines. Flow cytometric Cell cycle analysis showed that cells were arrested at the G1 phase after AKBA treatment and analysis showed that cyclin D1 and E, CDK 2 and 4 and phosphorylated Rb were reduces in AKBA-treated cells, while p21 expression was increases.<sup>26</sup> The growth inhibitory effect of AKBA was dependent on p21 of the apoptotic effect of AKBA, suggesting that p21 may be have protected cells against apoptosis by inducing a G1 arrest. AKBA inhibits cell growth in colon cancer cell.<sup>27</sup> Streffer et al. A clinical study with brain tumor patients was also performed, in which BSE was administered to 29

patients having glioma in three groups at different doses before surgery. After seven days of treatment, the reduction in the size of perifocal edema was found to be the greatest in the case of the group with the highest consumption of extract, to a lesser extent in the group used, without effect were observed doses lower than group receiving dose.<sup>28</sup> Sinha and associates. also found that boswellic acid also inhibited basic fibroblast growth factor [bFGF]-induce angiogenesis using the Matrigel Plug in vivo assay. Recent studies have also demonstrated that BAs can act as anti-angiogenic agents.<sup>29</sup> Singh et al. studied that BAs could even be considered as an alternative drugs to corticosteroids, as they have been shown to reduce cerebral peri-tumoural edema by modulating the of function P-glycoprotein [PgP]. Pgp has gained importance as the transporter, mainly for a drug disposition and the resulting clinical response; both BA and BSE inhibited the transport activity of PgP in the micromolecular range.<sup>30</sup> In the normal cells, ABA does not show apoptosis, ABA induces DNA fragmentation in melanoma and fibrosarcoma, ABA is a cytostatic rather than a cytotoxic agent, As it induces differentiation, apoptosis and cytostasis in various cell line, and can be used in a chemoprophylaxis intervention strategies, either to interrupt the appearance of a primary tumour or to reduced the probability of metastasis.<sup>31</sup> BAs has been shown to induce cell differentiation and inhibit topoisomerase I and II.<sup>32</sup>

### 3. Conclusion

Medicinal plants have a long history of therapeutic uses around the world and are still an important traditional medicinal plants and herbal product are safe for patients. Natural product has served as a primary source of medicines in the century, and about half of the pharmaceuticals used today are derivatives of natural product. Boswellic acid is made up of a chain of triterpene molecules cyclic molecules of which is produced by plants *Boswellia serrata*. Potential application of boswellic acid for cancer treatment have been focused here. In this ethnopharmacological review the anticancer potential of boswellic acid is highlighted. it may help the researchers for further investigation.

### 4. Source of Funding

None.

### 5. Conflict of Interest

None.

### 6. Acknowledgement

The Authors PG thanks the Author, and their UG Supervisor AGN for his genuine support. The Author AGN thanks Hon. Shri R.S. Yadav, President, Siddhant College of Pharmacy for his due support.

### References

- Petrovska BB. Historical review of medicinal plants' usage. *Pharmacognosy Rev.* 2012;6(11):1–5.
- Clark AM. Natural products as a resource for new drugs. *Pharmaceutical Res.* 1996;13(8):1133–41.
- Garg P, Deep A. Anti-cancer potential of boswellic acid: a mini. *J Drugs Med.* 2015;7(2):18–27.
- Hussain H, Al-Harrasi A, Csuk R, Shamraiz U, Green IR. Therapeutic potential of boswellic acids: a patent review. *Expert Opin Ther Patents.* 1990;27(1):81–90.
- Brendler T, Brinckmann JA, Schippmann U. Sustainable supply, a foundation for natural product development: The case of Indian frankincense (*Boswellia serrata* Roxb. ex Colebr.). *J Ethnopharmacology.* 2018;225(28):279–86.
- Paul M, Brüning G, Bergmann J, Jauch J. A Thin-layer Chromatography Method for the Identification of Three Different Olibanum Resins (*Boswellia serrata*, *Boswellia papyrifera* and *Boswellia carterii*, respectively, *Boswellia sacra*). *Phytochemical Anal.* 2012;23(2):184–9.
- Ilvessalo-Pfäffli MS. Botanical Classification. In: Fiber Atlas. Springer; 1995. p. 1. Available from: <https://link.springer.com/book/10.1007/978-3-662-07212-7>.
- Siddiqui MZ. *Boswellia serrata*, a potential antiinflammatory agent: an overview. *Indian J Pharm Sci.* 2011;73(3):255–61.
- Falzone L, Salomone S, Libra M. Evolution of cancer pharmacological treatments at the turn of the third millennium. *Front Pharmacol.* 2018;9:1300.
- Shah BA, Qazi GN, Taneja SC. Boswellic acids: a group of medicinally important compounds. *Natural Product Rep.* 2009;26(1):72–89.
- Kumar D, Kumar V, Jalwal P. Boswellic Acid-Potential tumors suppressant terpenoid-Photochemistry, Extraction and Isolation Methods-A comprehensive review study. *J Pharmacogn Phytochemistry.* 2016;5(3):231–9.
- Garg P, Deep A. Anti-cancer potential of boswellic acid: a mini. *J Drugs Med.* 2015;7(2):18–27.
- Syrovets T, Gschwend JE, Büchele B, Laumonier Y, Zugmaier W, Genze F. Inhibition of IκB kinase activity by acetyl-boswellic acids promotes apoptosis in androgen-independent PC-3 prostate cancer cells in vitro and in vivo. *J Biol Chem.* 2005;280(7):6170–80.
- Khan MA, Ali R, Parveen R, Najmi AK, Ahmad S. Pharmacological evidences for cytotoxic and antitumor properties of Boswellic acids from *Boswellia serrata*. *J Ethnopharmacol.* 2016;191:315–23. doi:10.1016/j.jep.2016.06.053.
- Rashan L, Hakkim FL, Idrees M, Essa M, Velusamy T, Al-Baloshi M. *Boswellia* gum resin and essential oils: Potential health benefits– An evidence based review. *Pharmacol Neurol Dis.* 2019;9(2):53–71.
- Tošić NG, Nikolić VD, Miljković VM, Nikolić LB. *Boswellia serrata*'resin isolates: Chemical composition and pharmacological activities. *Adv Technol.* 2022;11(1):76–87.
- Uthaman S, Snima KS, Annapoorna M, Ravindranath KC, Nair SV, Lakshmanan VK. Novel boswellic acids nanoparticles induces cell death in prostate cancer cells. *J Nat Prod.* 2012;5:100–8.
- Nandan CD, Reshmi P, Uthaman S, Snima KS, Unni AK, Kamath CR. Therapeutic properties of boswellic acid nanoparticles in prostate tumor-bearing BALB/c mice model. *J Nanopharmaceutics Drug Deliv.* 2013;1(1):30–7.
- Suhail MM, Wu W, Cao A, Mondalek FG, Fung KM, Shih PT. *Boswellia sacra* essential oil induces tumor cell-specific apoptosis and suppresses tumor aggressiveness in cultured human breast cancer cells. *BMC Complement Altern Med.* 2011;11(1):1–14.
- Ni X, Suhail MM, Yang Q, Cao A, Fung KM, Postier RG. Frankincense essential oil prepared from hydrodistillation of *Boswellia sacra* gum resins induces human pancreatic cancer cell death in cultures and in a xenograft murine model. *BMC Complement Altern Med.* 2012;12(1):1–14.
- Subramani R, Lakshmanaswamy R. Complementary and alternative medicine and breast cancer. . *Progress Mol Biol Transl Sci.*

- 2017;151:231–74.
22. Al-Harrasi A, Saidi S. Phytochemical analysis of the essential oil from botanically certified oleogum resin of *Boswellia sacra* (Omani Luban). *Molecules*. 2008;13(9):2181–9.
23. El-Din AN, Ata K, Abdel-Gawad AR, Fahmy NF. Impact of High Serum Levels of MMP-7, MMP-9, TGF- $\beta$  and PDGF Macrophage Activation Markers on Severity of COVID-19 in Obese-Diabetic Patients. *Infect Drug Resist*. 2021;14:4015–25.
24. Ahmed HH, Abdel-Rahman M, Salem FEH, Shalby AB, Lokman MS. Antitumor efficacy of *Boswellia serrata* extract in management of colon cancer induced in experimental animal. *Int J Pharm Pharm Sci*. 2013;5(3):379–89.
25. Harris DM, Go VLW. How dietary components protect from cancer. *Nut Cancer Preven*. 2006;4(1):27–58.
26. Liu JJ, Nilsson Å, Oredsson S, Badmaev V, Zhao WZ, Duan RD. Boswellic acids trigger apoptosis via a pathway dependent on caspase-8 activation but independent on Fas/Fas ligand interaction in colon cancer HT-29 cells. *Carcinogenesis*. 2002;23(12):2087–93.
27. Lv M, Zhuang X, Zhang Q, Cheng Y, Wu D, Wang X, et al. Acetyl-11-keto- $\beta$ -boswellic acid enhances the cisplatin sensitivity of non-small cell lung cancer cells through cell cycle arrest, apoptosis induction, and autophagy suppression via p21-dependent signaling pathway. *Cell Biol Toxicol*. 2021;37(2):209–28.
28. Streffer JR, Bitzer M, Schabet M, Dichgans J, Weller M. Response of radiochemotherapy-associated cerebral edema to a phytotherapeutic agent. *Neurol*. 2001;56(9):1219–21.
29. Sinha VR, Singh A, Singh S, Bhinge JR. Compression coated systems for colonic delivery of 5-fluorouracil. *J Pharm Pharmacol*. 2007;59(3):359–65.
30. Sharma A, Chhikara S, Ghodekar S, Bhatia S, Kharya M, Gajbhiye V. Phytochemical and Pharmacological investigations on *Boswellia serrata*. *Pharmacogn Rev*. 2009;3(5):206–15.
31. Garg P, Deep A. Anti-cancer potential of boswellic acid: a mini. *J Drugs Med*. 2015;7(2):18–27.
32. Tsutsui K, Tsutsui K, Sano K, Kikuchi A, Tokunaga A. Involvement of DNA topoisomerase II $\beta$  in neuronal differentiation. *J Biol Chem*. 2001;276(8):5769–78.

### Author biography

**Amit Gajanan Nerkar**, Professor and Research Head  
 <https://orcid.org/0000-0002-1377-8466>

**Pallavi Gade**, Student

**Cite this article:** Nerkar AG, Gade P. Ethnopharmacological review of *boswellia serrata* for anticancer activity. *Curr Trends Pharm Pharm Chem* 2022;4(4):144-147.