



Original Research Article

***In Vitro* anticancer screening of herbal syrup formulation of allium cepa extract**Amit G. Nerkar^{1,*}, Ashutosh Pansare¹¹Dept. of Pharmacy, CAYMET's Siddhant College of Pharmacy, Sudumbare, Pune, Maharashtra, India

ARTICLE INFO

Article history:

Received 14-09-2023

Accepted 30-10-2023

Available online 06-11-2023

Keywords:

Herbal Formulation

Syrup

Onion

Allium cepa

in vitro

Anticancer screening

ABSTRACT

The formulation of herbal syrup of extract of Allium Cepa was reported in the previous paper. The herbal syrup of extract was evaluated here for *in vitro* anticancer activity. The *in vitro* anticancer activity was evaluated against 2 cell lines namely human lung cancer cell line A-549 and human breast cancer cell line MCF7. The results indicate that there is no significant anticancer activity of the herbal syrup formulation of the extract as compared with the standard Adriamycin.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

1.1. Role of allium cepa

Allium Cepa or the onion has significant role in day-to-day life. Mostly it is consumed as vegetable fruit, spices, and adds delicious flavour to the food.¹ Allium Cepa is well known for its antidiabetic properties. Also, it has been reported to have aphrodisiac along with Safed Musli,² blood purifying³ and as a fibre to the diet.⁴ The well know utilities of onion or Allium Cepa can be found in Ayurveda. Thus, it can be consumed as extract and as a nutraceutical for complementary medicine and naturopathic purposes.⁵ Further it has been reported to have anticancer properties.^{6–8}

1.2. Medicated herbal syrup of onion or allium cepa extract

The formulation and evaluation or optimization of formulation of herbal syrup of extract of Allium Cepa was reported by Nerkar et al.⁹ in agreement of the standard guidelines of Indian Pharmacopoeia i.e., the guidelines

to optimize the syrup dosage form. Thus, in this paper we report here the *in vitro* anticancer screening of the formulation of Allium Cepa or Onion against A549 Human Lung Cancer Cell Line and MCF7 Human Breast cancer Cell Line.

2. Materials and Methods

The syrup was dried in petri dishes and 0.5g dried syrup was dissolved in 5 ml of water to give a concentration on 100mg/ ml. Further anticancer cytotoxicity assay was performed on 2 Cell lines viz Human Lung Cancer Cell line A549 and Human Breast cancer Cell Line MCF7. The dried formulation of Allium Cepa extract was coded as APO.

2.1. *In vitro* anticancer activity against cancer cell lines (anticancer cytotoxicity assay)

2.1.1. SRB assay

The assay relies on the ability of SRB to bind to protein components of cells that have been fixed to tissue-culture plates by trichloroacetic acid (TCA). The cell lines were grown in RPMI 1640 medium containing 10% fetal bovine

* Corresponding author.

E-mail address: dragnerkar@gmail.com (A. G. Nerkar).

Table 1: *In vitro* anticancer assay of herbal syrup of allium cepa (APO) against A549

Human Lung Cancer Cell Line A549																
% Control Growth																
Drug Concentrations ($\mu\text{g/ml}$)																
Experiment 1				Experiment 2				Experiment 3				Average Values				
10	20	40	80	10	20	40	80	10	20	40	80	10	20	40	80	
APO	47.4	32.8	26.7	20.6	40.5	28.1	25.9	20.0	41.4	27.6	18.2	16.3	43.1	29.5	23.6	19.0
ADR	3.7	3.0	0.2	-1.2	4.4	3.4	0.5	-4.6	2.1	2.1	1.2	0.3	3.4	2.8	0.6	-1.9

Table 2: *In vitro* anticancer assay of herbal syrup of allium cepa (APO) against MCF7

Human Breast Cancer Cell Line MCF7																
% Control Growth																
Drug Concentrations ($\mu\text{g/ml}$)																
Experiment 1				Experiment 2				Experiment 3				Average Values				
10	20	40	80	10	20	40	80	10	20	40	80	10	20	40	80	
APO	49.0	47.4	40.0	23.6	56.2	42.6	37.8	33.3	61.8	43.8	35.9	25.9	55.7	44.6	37.9	27.6
ADR	3.7	3.0	0.2	-1.2	4.4	3.4	0.5	-4.6	2.1	2.1	1.2	0.3	3.4	2.8	0.6	-1.9

Table 3: *In vitro* anticancer assay showing LC50, TGI and GI50 of APO against A549

Drug concentrations ($\mu\text{g/ml}$) calculated from graph				
MCF7	LC50		TGI	GI50
APO	>80		>80	21.6
ADR	3.7		3.0	0.2

Table 4: *In vitro* anticancer assay showing LC50, TGI and GI50 of APO against MCF7

Drug concentrations ($\mu\text{g/ml}$) calculated from graph				
MCF7	LC50		TGI	GI50
APO	>80		>80	35.5
ADR	3.7		3.0	0.2

serum and 2 mM L-glutamine. For present screening experiment, cells were inoculated into 96 well microtiter plates in 100 μL at plating densities. After cell inoculation, the microtiter plates were incubated at 37°C, 5% CO₂, 95% air and 100% relative humidity for 24 h prior to addition of experimental drugs. Experimental drugs were initially solubilized in dimethyl sulfoxide at 100mg/ml and diluted to 1mg/ml using water and stored frozen prior to use. At the time of drug addition, an aliquot of frozen concentrate (1mg/ml) was thawed and diluted to 100 $\mu\text{g/ml}$, 200 $\mu\text{g/ml}$, 400 $\mu\text{g/ml}$ and 800 $\mu\text{g/ml}$ with complete medium containing test article. Aliquots of 10 μl of these different drug dilutions were added to the appropriate microtiter wells already containing 90 μl of medium, resulting in the required final drug concentrations i.e. 10 $\mu\text{g/ml}$, 20 $\mu\text{g/ml}$, 40 $\mu\text{g/ml}$, 80 $\mu\text{g/ml}$. After compound addition, plates were incubated at standard conditions for 48 hours and assay was terminated by the addition of cold TCA. Cells were fixed in situ by the gentle addition of 50 μl of cold 30% (w/v) TCA (final concentration, 10% TCA) and incubated for 60 minutes at 4°C. The supernatant was discarded; the plates were washed five times with tap water and air dried. Sulforhodamine B (SRB) solution (50 μl) at 0.4% (w/v) in

1% acetic acid was added to each of the wells, and plates were incubated for 20 minutes at room temperature. After staining, unbound dye was recovered and the residual dye was removed by washing five times with 1% acetic acid. The plates were air dried. Bound stain was subsequently eluted with 10mM trizma base, and the absorbance was read on a plate reader at a wavelength of 540nm with 690nm reference wavelength. Percent growth was calculated on a plate-by-plate basis for test wells relative to control wells. Percent Growth was expressed as the ratio of average absorbance of the test well to the average absorbance of the control wells x 100. Using the six absorbance measurements [time zero (Tz), control growth (C), and test growth in the presence of drug at the four concentration levels (Ti)], the percentage growth was calculated at each of the drug concentration levels. Percentage growth inhibition was calculated as: $[\text{Ti}/\text{C}] \times 100\%$ Percentage growth inhibition, total growth inhibition TGI) and LC50 was calculated. GI50 value of $\leq 10 \mu\text{g/ml}$ is considered to demonstrate activity in case of pure compounds. For extracts, GI50 value $\leq 20 \mu\text{g/ml}$ is considered to demonstrate activity. Above three parameters were calculated only when the level of activity was observed. The values were expressed as greater or less

than maximum or minimum concentration tested when the effect was not reached or exceeded.^{10,11}

3. Result and Discussion

From results as mentioned in Tables 1, 2, 3 and 4., it is exemplified that the herbal syrup formulation of Allium Cepa or Onion extract does not show any significant anticancer activity at concentrations of 10µg/ml, 20µg/ml, 40µg/ml and 80µg/ml as compared to parameters of LC50, TGI and GI50 and standard drug Adriamycin used for the assay. However, there is a possibility that it may show anticancer activity against other cancer cell lines which is a part of further research.

4. Conclusion

The herbal formulation of syrup of Allium Cepa extract does not show any significant anticancer activity against Human A549 and Human MCF-7 Cell Line.

5. Source of Funding

None.

6. Conflict of Interest

None.


7. Acknowledgment

The Authors AP thanks UG Supervisor AGN for his genuine support.

References

1. Kurup AH, Deotale S, Rawson A, Patras A. Thermal Processing of Herbs and Spices. . In: and others, editor. Herbs, Spices and Medicinal Plants: Processing, Health Benefits and Safety. vol. 12; 2020. p. 1–21. doi:10.1002/9781119036685..
2. Bhat MH, Fayaz M, Kumar A, Jain AK. Phytochemical, pharmacological and nutritional profile of Chlorophytum tuberosum (Roxb.) Baker (Safed musli): A review. *Int J Theor Appl Sci.* 2018;10(1):93–102.
3. Pramanik K, Tripathy P, Mandal P, Pradhan M, Biswal M. Effect of micronutrients on quality of onion (*Allium cepa* L.). *Int J Chem Stud.* 2018;6(6):1324–31.
4. Mahmood N, Muazzam MA, Ahmad M, Hussain S, Javed W. Phytochemistry of Allium cepa L.(Onion): An overview of its nutritional and pharmacological importance. . *Sci Inquiry Rev.* 2021;5:41–59.
5. Pizzorno JE, Murray MT. Textbook of natural medicine. 5th ed. Seattle, WA; 1999.
6. Abdelrahman M, Mahmoud HY, El-Sayed M, Tanaka S, Tran LS. Isolation and characterization of Cepa2, a natural alliospiroside A, from shallot (*Allium cepa* L. Aggregatum group) with anticancer activity. *Plant Physiol Biochem.* 2017;116:167–73.
7. Kumar M, Barbhai MD, Hasan M, Punia S, Dhupal S, Rais N, et al. Onion (*Allium cepa* L.) peels: A review on bioactive compounds and biomedical activities. *Biomed Pharmacother.* 2022;146:112498. doi:10.1016/j.biopha.2021.112498.
8. Fadholly A, Ansori AN, Jayanti S, Proboningrat A, Kusala MK, Putri N, et al. Cytotoxic effect of Allium cepa L. extract on human colon cancer (WiDr) cells: in vitro study. *Res J Pharm Technol.* 2019;12(7):3483–9.
9. Nerkar AG, Pansare A. Formulation and evaluation of herbal syrup of onion (*Allium Cepa*). *Curr Trends Pharmacy Pharm Chem.* 2023;5(3):97–100.
10. Skehn P, Storeng R, Scudiero A, Monks J, Mcmohan D, Vistica D, et al. New colorimetric cytotoxicity assay for anticancer drug screening. *J National Cancer Institute.* 1990;82(13):1107–12.
11. Vichai V, Kirtikara K. Sulforhodamine B colorimetric assay for cytotoxicity drug screening. *Nature Protocols.* 2006;1(3):1112–6.

Author biography

Amit G. Nerkar, Professor and Research Head  <https://orcid.org/0000-0002-1377-8466>

Ashutosh Pansare, Student

Cite this article: Nerkar AG, Pansare A. *In Vitro* anticancer screening of herbal syrup formulation of allium cepa extract. *Curr Trends Pharm Pharm Chem* 2023;5(4):133-135.